

# Melnikov Function associated to Morse Real Polynomials

M.Uribe \*

August 9, 2013

*Departamento de Matemática y Física Aplicadas,  
Facultad de Ingeniería  
Universidad Católica de la Sma. Concepción  
Casilla 297, Concepción, Chile.*

## Abstract

It is known that the Principal Melnikov Function is generically an Abelian integral. We give a sufficient condition on monodromy to ensure that it is an Abelian Integral also in Non generic cases.

In non generic cases it is an iterated integral [1]. In [2], [3], the author gives a precise description of the Principal Melnikov Function, an iterated integral of length at most 2, involving a logarithmic function with only one ramification at a point at infinity. We extend this result to some isomonodromic families of real Morse polynomial [4].

**Keywords:** Abelian integrals, Melnikov functions, First Return Map, Monodromy.

## References

- [1] Gavrilov, L. *Higher order Poincaré-Pontryagin functions and iterated path integrals*. Ann. Fac. Sci. Toulouse Math. (6) 14 (2005), no. 4, 663–682.
- [2] Uribe M. *Principal Poincaré-Pontryagin Function of Polynomial Perturbations of the Hamiltonian Triangle*. Journal of Dynamical and Control Systems, Vol 12, 1 (2006), 109-134.
- [3] Uribe, M. *Principal Poincaré-Pontryagin function associated to polynomial perturbations of a product of  $(d + 1)$  straight lines*. J. Differential Equations 246 (2009), no. 4, 1313–1341.
- [4] Pelletier, M. and Uribe, M. *The displacement map associated to some families of Morse real polynomials*. Submitted to Nonlinearity (2013).

---

\*Trabajo parcialmente financiado por Proyecto Fondecyt11080250, E-mail address: muribe@ucsc.cl